

General Certificate of Education

Physics 5451

Specification A

PA01 Particles, Radiation and Quantum Phenomena

Mark Scheme

2007 examination - June series

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available to download from the AQA Website: www.aqa.org.uk

Copyright © 2007 AQA and its licensors. All rights reserved.

COPYRIGHT

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

The Assessment and Qualifications Alliance (AQA) is a company limited by guarantee registered in England and Wales (company number 3644723) and a registered charity (registered charity number 1073334). Registered address: AQA, Devas Street, Manchester M15 6EX Dr Michael Cresswell Director General

Instructions to Examiners

- 1 Give due credit to alternative treatments which are correct. Give marks for what is correct; do not deduct marks because the attempt falls short of some ideal answer. Where marks are to be deducted for particular errors specific instructions are given in the marking scheme.
- 2 Do not deduct marks for poor written communication. Refer the script to the Awards meeting if poor presentation forbids a proper assessment. In each paper candidates may be awarded up to two marks for the Quality of Written Communication in cases of required explanation or description. Use the following criteria to award marks:
 - 2 marks: Candidates write legibly with accurate spelling, grammar and punctuation; the answer containing information that bears some relevance to the question and being organised clearly and coherently. The vocabulary should be appropriate to the topic being examined.
 - 1 mark: Candidates write with reasonably accurate spelling, grammar and punctuation; the answer containing some information that bears some relevance to the question and being reasonably well organised. Some of the vocabulary should be appropriate to the topic being examined.
 - 0 marks: Candidates who fail to reach the threshold for the award of one mark.
- 3 An arithmetical error in an answer should be marked AE thus causing the candidate to lose one mark. The candidate's incorrect value should be carried through all subsequent calculations for the question and, if there are no subsequent errors, the candidate can score all remaining marks (indicated by ticks). These subsequent ticks should be marked CE (consequential error).
- 4 With regard to incorrect use of significant figures, normally two, three or four significant figures will be acceptable. Exceptions to this rule occur if the data in the question is given to, for example, five significant figures as in values of wavelength or frequency in questions dealing with the Doppler effect, or in atomic data. In these cases up to two further significant figures will be acceptable. The maximum penalty for an error in significant figures is **one mark per paper**. When the penalty is imposed, indicate the error in the script by SF and, in addition, write SF opposite the mark for that question on the front cover of the paper to obviate imposing the penalty more than once per paper.
- 5 No penalties should be imposed for incorrect or omitted units at intermediate stages in a calculation or which are contained in brackets in the marking scheme. Penalties for unit errors (incorrect or omitted units) are imposed only at the stage when the final answer to a calculation is considered. The maximum penalty is **one mark per question**.
- 6 All other procedures, including the entering of marks, transferring marks to the front cover and referrals of scripts (other than those mentioned above) will be clarified at the standardising meeting of examiners.

Question 1		
(a)	neutrons (= 55 – 26) = 29 ✓	2
	electrons = 26 protons = 26 \checkmark	2
(b)	charge = 26 × 1.6 × 10 ⁻¹⁹ ✓	
	mass = 55 × 1.67 × 10 ⁻²⁷ ✓	3
	charge/mass = $4.5 \times 10^7 \mathrm{C kg^{-1}}$	
(C)	a = 55 ✓	2
	b = 27 ✓	2
	Total	7

PA01 Particles, Radiation and Quantum Phenomena

Question 2		
(a)	qqq (or <u>qqq</u>) ✓	1
(b)	meson ✓	2
	qq̄✓	L
(C)	neutral pion or $\pi^{\circ} \checkmark$	1
(d)	proton 🗸	1
(e)	as strangeness is not conserved \checkmark	
	or weak link interaction is the only interaction that changes the type of quark \checkmark	1
	Total	6

Question 3		
(a)	photons have definite wavelengths/line spectra/frequency \checkmark	
	so photons have discrete energies \checkmark	
	photons are emitted when electrons/atoms move down from one level to another \checkmark	max 3
	the energy gaps between energy levels are fixed \checkmark	
(b) (i)	$E_4 = -1.4 \times 10^{-19} (J) \checkmark$	
	use of $hf = E_1 - E_2 \checkmark$	
	$hf = (-1.4 \times 10^{-19}5.5 \times 10^{-19}) = 4.1 \times 10^{-19} \text{ J} \checkmark$	4
(ii)	$(f = 4.1 \times 10^{-19}/6.6 \times 10^{-34})$	
	= 6.2×10^{14} Hz \checkmark [possible CE from (i)]	
(c)	6 ✓	1
	Total	8

Question 4		
(a) (i)	(otherwise) α particles may be absorbed/so α particles can penetrate the foil \checkmark α particles must only be scattered once \checkmark	max 3
(ii)	to have definite location where the scattering takes place \checkmark	
	so the scattering angle can be determined accurately \checkmark	
(b)	backscattering/scattering angles greater than 90° \checkmark	
	occurs because α particles must collide with a target (much) momentum massive then themselves \checkmark	re 2
(C)	top ray – showing at least 10° deflection upwards \checkmark	
	middle ray – approaches closer than about 1 cm but not touching nucleus and is backscattered ✓	3
	bottom ray – smooth curve deflected downwards at an angle greater than the top ray is deflected upwards \checkmark	
	Tc	otal 8

Que	stion 5			
(a)		minimum energy ✓	2	
		required to remove an electron from a metal \checkmark	2	
(b)	(i)	(using $E = hf$ and $c = f\lambda$)		
		$\lambda = hc/E \checkmark$		
		= $6.6 \times 10^{-34} \times 3.0 \times 10^{8} / 7.9 \times 10^{-19}$ \checkmark		
		= 2.5×10^{-7} m \checkmark		
	(ii)	(use of $hf = \phi + E_k$)	6	
		$\phi = (7.9 \times 10^{-19} - 4.2 \times 10^{-19}) \checkmark = 3.7 \times 10^{-19} \text{ (J) }\checkmark$		
	(iii)	$\phi = 3.7 \times 10^{-19} / 1.6 \times 10^{-19}$		
		= 2.3 (eV) ✓ [possible CE from (ii)]		
(C)		the number of photoelectrons emitted per second will be increased ✓	mov 3	
		as the number of incident photons per second is increased \checkmark		
		but the maximum kinetic energy of the photoelectrons will remain constant \checkmark	max 5	
		a photon gives up all its energy in one collision \checkmark		
		Total	11	

Que	stion 6		
(a)		the angle to the normal inside the material \checkmark	
		which results in 90⁰ refraction in air ✓	2
		(definitions in terms of limits of TIR or limits of refraction are also allowed)	
(b)	(i)	use of $n = 1 / \sin \theta_c$	
		$\theta_{\rm c} \ (= \sin^{-1} 1 \ / \ 1.6) = 39^{\circ} \checkmark$	
	(ii)	TIR on hypotenuse and angles drawn equal \checkmark	
		refraction on top surface ✓	6
	(iii)	use of $_1n_2 = \sin \theta_1 / \sin \theta_2$ and $_1n_2 = n_2 / n_1 \checkmark$	
		$n_2 = 1.6 \times \sin 60 / \sin 90 \checkmark$	
		= 1.4 🗸	
		Total	8

Quality of Written Communication: Q3 (a) and/or Q5 (c)	2
--	---