

**PHYSICS (SPECIFICATION A)**  
**Unit 4 Waves, Fields and Nuclear Energy**

**PA04**

**Section A**

Thursday 16 June 2005 Morning Session

**In addition to this paper you will require:**

- an objective test answer sheet;
- a black ball-point pen;
- a calculator;
- a question paper/answer book for Section B (enclosed).

Time allowed: The total time for Section A and Section B of this paper is 1 hour 30 minutes

**Instructions**

- Use a black ball-point pen. Do **not** use pencil.
- Answer **all** questions in this section.
- For each question there are four responses. When you have selected the response which you think is the most appropriate answer to a question, mark this response on your answer sheet.
- Mark all responses as instructed on your answer sheet. If you wish to change your answer to a question, follow the instructions on your answer sheet.
- Do all rough work in this book, **not** on the answer sheet.

**Information**

- The maximum mark for this Section is 30.
- Section A and Section B of this paper together carry 15% of the total marks for Physics Advanced.
- All questions in Section A carry equal marks. No deductions will be made for incorrect answers.
- A *Data Sheet* is provided on pages 3 and 4. You may wish to detach this perforated sheet at the start of the examination.
- The question paper/answer book for Section B is enclosed within this question paper.

**Data Sheet**

- A perforated *Data Sheet* is provided as pages 3 and 4 of this question paper.
- This sheet may be useful for answering some of the questions in the examination.
- You may wish to detach this sheet before you begin work.

**Fundamental constants and values**

Quantity	Symbol	Value	Units
speed of light in vacuo	$c$	$3.00 \times 10^8$	$\text{m s}^{-1}$
permeability of free space	$\mu_0$	$4\pi \times 10^{-7}$	$\text{H m}^{-1}$
permittivity of free space	$\epsilon_0$	$8.85 \times 10^{-12}$	$\text{F m}^{-1}$
charge of electron	$e$	$1.60 \times 10^{-19}$	C
the Planck constant	$h$	$6.63 \times 10^{-34}$	J s
gravitational constant	$G$	$6.67 \times 10^{-11}$	$\text{N m}^2 \text{kg}^{-2}$
the Avogadro constant	$N_A$	$6.02 \times 10^{23}$	$\text{mol}^{-1}$
molar gas constant	$R$	8.31	$\text{J K}^{-1} \text{mol}^{-1}$
the Boltzmann constant	$k$	$1.38 \times 10^{-23}$	$\text{J K}^{-1}$
the Stefan constant	$\sigma$	$5.67 \times 10^{-8}$	$\text{W m}^{-2} \text{K}^{-4}$
the Wien constant	$\alpha$	$2.90 \times 10^{-3}$	m K
electron rest mass	$m_e$	$9.11 \times 10^{-31}$	kg
(equivalent to $5.5 \times 10^{-4}u$ )			
electron charge/mass ratio	$em_e$	$1.76 \times 10^{11}$	$\text{C kg}^{-1}$
proton rest mass	$m_p$	$1.67 \times 10^{-27}$	kg
(equivalent to 1.00728u)			
proton charge/mass ratio	$em_p$	$9.58 \times 10^7$	$\text{C kg}^{-1}$
neutron rest mass	$m_n$	$1.67 \times 10^{-27}$	kg
(equivalent to 1.00867u)			
gravitational field strength	$g$	9.81	$\text{N kg}^{-1}$
acceleration due to gravity	$g$	9.81	$\text{m s}^{-2}$
atomic mass unit	$u$	$1.661 \times 10^{-27}$	kg
(1u is equivalent to 931.3 MeV)			

**Fundamental particles**

Class	Name	Symbol	Rest energy /MeV
photon	photon	$\gamma$	0
lepton	neutrino	$\nu_e$	0
		$\nu_\mu$	0
	electron	$e^\pm$	0.510999
	muon	$\mu^\pm$	105.659
mesons	pion	$\pi^\pm$	139.576
		$\pi^0$	134.972
	kaon	$K^\pm$	493.821
		$K^0$	497.762
baryons	proton	p	938.257
	neutron	n	939.551

**Properties of quarks**

Type	Charge	Baryon number	Strangeness
u	$+\frac{2}{3}$	$+\frac{1}{3}$	0
d	$-\frac{1}{3}$	$+\frac{1}{3}$	0
s	$-\frac{1}{3}$	$+\frac{1}{3}$	-1

**Geometrical equations**

- arc length =  $r\theta$
- circumference of circle =  $2\pi r$
- area of circle =  $\pi r^2$
- area of cylinder =  $2\pi rh$
- volume of cylinder =  $\pi r^2 h$
- area of sphere =  $4\pi r^2$
- volume of sphere =  $\frac{4}{3}\pi r^3$

**Mechanics and Applied Physics**

- $v = u + at$
- $s = \left(\frac{u+v}{2}\right)t$
- $s = ut + \frac{at^2}{2}$
- $v^2 = u^2 + 2as$
- $F = \frac{\Delta(mv)}{\Delta t}$
- $P = Fv$
- efficiency =  $\frac{\text{power output}}{\text{power input}}$
- $\omega = \frac{v}{r} = 2\pi f$
- $a = \frac{v^2}{r} = r\omega^2$
- $I = \sum mr^2$
- $E_k = \frac{1}{2}I\omega^2$
- $\omega_2 = \omega_1 + at$
- $\theta = \omega_1 t + \frac{1}{2}at^2$
- $\omega_2^2 = \omega_1^2 + 2a\theta$
- $\theta = \frac{1}{2}(\omega_1 + \omega_2)t$
- $T = I\alpha$
- angular momentum =  $I\omega$
- $W = T\theta$
- $P = T\omega$
- angular impulse = change of angular momentum =  $Tt$
- $\Delta Q = \Delta U + \Delta W$
- $\Delta W = p\Delta V$
- $pV^\gamma = \text{constant}$
- work done per cycle = area of loop
- input power = calorific value  $\times$  fuel flow rate
- indicated power as (area of  $p-V$  loop)  $\times$  (no. of cycles/s)  $\times$  (no. of cylinders)
- friction power = indicated power - brake power
- efficiency =  $\frac{W}{Q_{in}} = \frac{Q_{in} - Q_{out}}{Q_{in}}$
- maximum possible efficiency =  $\frac{T_H - T_C}{T_H}$

**Fields, Waves, Quantum Phenomena**

- $g = \frac{F}{m}$
- $g = -\frac{GM}{r^2}$
- $g = -\frac{\Delta V}{\Delta x}$
- $V = -\frac{GM}{r}$
- $a = -(2\pi f)^2 x$
- $v = \pm 2\pi f \sqrt{A^2 - x^2}$
- $x = A \cos 2\pi ft$
- $T = 2\pi \sqrt{\frac{m}{k}}$
- $T = 2\pi \sqrt{\frac{l}{g}}$
- $\lambda = \frac{\omega S}{D}$
- $d \sin \theta = n\lambda$
- $\theta \approx \frac{\lambda}{D}$
- ${}_1n_2 = \frac{\sin \theta_1}{\sin \theta_2} = \frac{c_1}{c_2}$
- ${}_1n_2 = \frac{n_2}{n_1}$
- $\sin \theta_c = \frac{1}{n}$
- $E = hf$
- $hf = \phi + E_k$
- $hf = E_1 - E_2$
- $\lambda = \frac{h}{p} = \frac{h}{mv}$
- $c = \frac{1}{\sqrt{\mu_0 \epsilon_0}}$

**Electricity**

- $\epsilon = \frac{E}{Q}$
- $\epsilon = I(R + r)$
- $\frac{1}{R_T} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots$
- $R_T = R_1 + R_2 + R_3 + \dots$
- $P = I^2 R$
- $E = \frac{F}{Q} = \frac{V}{d}$
- $E = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2}$
- $E = \frac{1}{2} QV$
- $F = BI$
- $F = BQv$
- $Q = Q_0 e^{-t/RC}$
- $\Phi = BA$

Turn over

$$\text{magnitude of induced e.m.f.} = N \frac{\Delta\Phi}{\Delta t}$$

$$I_{\text{rms}} = \frac{I_0}{\sqrt{2}}$$

$$V_{\text{rms}} = \frac{V_0}{\sqrt{2}}$$

### Mechanical and Thermal Properties

$$\text{the Young modulus} = \frac{\text{tensile stress}}{\text{tensile strain}} = \frac{F}{A} \frac{l}{e}$$

$$\text{energy stored} = \frac{1}{2} Fe$$

$$\Delta Q = mc \Delta\theta$$

$$\Delta Q = ml$$

$$pV = \frac{1}{3} Nmc^2$$

$$\frac{1}{2} mc^2 = \frac{3}{2} kT = \frac{3RT}{2N_A}$$

### Nuclear Physics and Turning Points in Physics

$$\text{force} = \frac{eV_p}{d}$$

$$\text{force} = Bev$$

$$\text{radius of curvature} = \frac{mv}{Be}$$

$$\frac{eV}{d} = mg$$

$$\text{work done} = eV$$

$$F = 6\pi\eta rv$$

$$I = k \frac{I_0}{x^2}$$

$$\frac{\Delta N}{\Delta t} = -\lambda N$$

$$\lambda = \frac{h}{\sqrt{2}meV}$$

$$N = N_0 e^{-\lambda t}$$

$$T_{\frac{1}{2}} = \frac{\ln 2}{\lambda}$$

$$R = r_0 A^{\frac{1}{3}}$$

$$E = mc^2 = \frac{m_0 c^2}{\left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}}$$

$$l = l_0 \left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}$$

$$t = \frac{t_0}{\left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}}$$

### Astrophysics and Medical Physics

Body	Mass/kg	Mean radius/m
Sun	$2.00 \times 10^{30}$	$7.00 \times 10^8$
Earth	$6.00 \times 10^{24}$	$6.40 \times 10^6$

1 astronomical unit =  $1.50 \times 10^{11}$  m

1 parsec = 206265 AU =  $3.08 \times 10^{16}$  m = 3.26 ly

1 light year =  $9.45 \times 10^{15}$  m

Hubble constant ( $H$ ) =  $65 \text{ km s}^{-1} \text{ Mpc}^{-1}$

$$M = \frac{\text{angle subtended by image at eye}}{\text{angle subtended by object at unaided eye}}$$

$$M = \frac{f_o}{f_e}$$

$$m - M = 5 \log \frac{d}{10}$$

$$\lambda_{\text{max}} T = \text{constant} = 0.0029 \text{ m K}$$

$$v = Hd$$

$$P = \sigma AT^4$$

$$\frac{\Delta f}{f} = \frac{v}{c}$$

$$\frac{\Delta \lambda}{\lambda} = -\frac{v}{c}$$

$$R_s \approx \frac{2GM}{c^2}$$

### Medical Physics

$$\text{power} = \frac{1}{f}$$

$$\frac{1}{u} + \frac{1}{v} = \frac{1}{f} \text{ and } m = \frac{v}{u}$$

$$\text{intensity level} = 10 \log \frac{I}{I_0}$$

$$I = I_0 e^{-\mu x}$$

$$\mu_m = \frac{\mu}{\rho}$$

### Electronics

Resistors

Preferred values for resistors (E24)

Series: 1.0 1.1 1.2 1.3 1.5 1.6 1.8 2.0 2.2 2.4 2.7 3.0 3.3 3.6 3.9 4.3 4.7 5.1 5.6 6.2 6.8 7.5 8.2 9.1 ohms

and multiples that are ten times greater

$$Z = \frac{V_{\text{rms}}}{I_{\text{rms}}}$$

$$\frac{1}{C_T} = \frac{1}{C_1} + \frac{1}{C_2} + \frac{1}{C_3} + \dots$$

$$C_T = C_1 + C_2 + C_3 + \dots$$

$$X_C = \frac{1}{2\pi f C}$$

### Alternating Currents

$$f = \frac{1}{T}$$

### Operational amplifier

$$G = \frac{V_{\text{out}}}{V_{\text{in}}} \quad \text{voltage gain}$$

$$G = -\frac{R_f}{R_1} \quad \text{inverting}$$

$$G = 1 + \frac{R_f}{R_1} \quad \text{non-inverting}$$


$$V_{\text{out}} = -R_f \left( \frac{V_1}{R_1} + \frac{V_2}{R_2} + \frac{V_3}{R_3} \right) \quad \text{summing}$$

## SECTION A

In this section each item consists of a question or an incomplete statement followed by four suggested answers or completions. You are to select the most appropriate answer in each case.

You are advised to spend approximately **30 minutes** on this section.

- 1 A spring is suspended from a fixed point. A mass attached to the spring is set into vertical undamped simple harmonic motion. When the mass is at its lowest position, which one of the following has its minimum value?
- A the potential energy of the system  
 B the kinetic energy of the mass  
 C the acceleration of the mass  
 D the tension in the spring
- 2 The time period of a simple pendulum is doubled when the length of the pendulum is increased by 3.0 m. What is the original length of the pendulum?
- A 1.0 m  
 B 1.5 m  
 C 3.0 m  
 D 6.0 m
- 3 The diagram shows a snapshot of a wave on a rope travelling from left to right.



At the instant shown, point **P** is at maximum displacement and point **Q** is at zero displacement. Which one of the following lines, **A** to **D**, in the table correctly describes the motion of **P** and **Q** in the next half-cycle?

	<b>P</b>	<b>Q</b>
<b>A</b>	falls then rises	rises
<b>B</b>	falls then rises	rises then falls
<b>C</b>	falls	falls
<b>D</b>	falls	rises then falls

Turn over ►

- 4 The speed of sound in water is  $1500 \text{ m s}^{-1}$ . For a sound wave in water having frequency  $2500 \text{ Hz}$ , what is the minimum distance between two points at which the vibrations are  $\frac{\pi}{3}$  rad out of phase?
- A 0.05 m  
B 0.10 m  
C 0.15 m  
D 0.20 m
- 5 Which one of the following properties of light waves do polarising sunglasses depend on for their action?
- Light waves may
- A interfere constructively.  
B interfere destructively.  
C be polarised when reflected from a surface.  
D be polarised by the lens in the eye.
- 6 Light of wavelength  $\lambda$  is incident normally on a diffraction grating for which adjacent lines are a distance  $3\lambda$  apart. What is the angle between the second order maximum and the straight-through position?
- A  $9.6^\circ$   
B  $20^\circ$   
C  $42^\circ$   
D There is no second order maximum.
- 7 The Earth has density  $\rho$  and radius  $R$ . The gravitational field strength at the surface is  $g$ . What is the gravitational field strength at the surface of a planet of density  $2\rho$  and radius  $2R$ ?
- A  $g$   
B  $2g$   
C  $4g$   
D  $16g$
- 8 A particle of mass  $m$  moves in a circle of radius  $r$  at uniform speed, taking time  $T$  for each revolution. What is the kinetic energy of the particle?
- A  $\frac{\pi^2 m r}{T^2}$   
B  $\frac{\pi^2 m r^2}{T^2}$   
C  $\frac{2\pi^2 m r^2}{T}$   
D  $\frac{2\pi^2 m r^2}{T^2}$

- 9 Two protons, each of mass  $m$  and charge  $e$ , are a distance  $d$  apart. Which one of the following expressions correctly gives the ratio  $\left(\frac{\text{electrostatic force}}{\text{gravitational force}}\right)$  for the forces acting between them?


A  $\frac{4\pi\epsilon_0 e^2}{Gm^2}$

B  $\frac{Ge^2}{4\pi\epsilon_0 m^2}$

C  $\frac{e^2 m^2}{4\pi\epsilon_0 G}$

D  $\frac{e^2}{4\pi\epsilon_0 Gm^2}$

- 10 The graph shows how the gravitational potential,  $V$ , varies with the distance,  $r$ , from the centre of the Earth.




What does the gradient of the graph at any point represent?

- A the magnitude of the gravitational field strength at that point  
B the magnitude of the gravitational constant  
C the mass of the Earth  
D the potential energy at the point where the gradient is measured

Turn over ►

11




The diagram shows two charges,  $+4\ \mu\text{C}$  and  $-16\ \mu\text{C}$ , 120 mm apart. What is the distance from the  $+4\ \mu\text{C}$  charge to the point between the two charges, where the resultant electric potential is zero?

- A 24 mm  
B 40 mm  
C 80 mm  
D 96 mm
- 12 An electron travelling at constant speed enters a uniform electric field at right angles to the field. While the electron is in the field it accelerates in a direction which is
- A in the same direction as the electric field.  
B in the opposite direction to the electric field.  
C in the same direction as the motion of the electron.  
D in the opposite direction to the motion of the electron.
- 13 A  $1000\ \mu\text{F}$  capacitor and a  $10\ \mu\text{F}$  capacitor are charged so that the potential difference across each of them is the same. The charge stored in the  $1000\ \mu\text{F}$  capacitor is  $Q_1$  and the charge stored in the  $10\ \mu\text{F}$  capacitor is  $Q_2$ .
- What is the ratio  $\frac{Q_1}{Q_2}$  ?
- A 100  
B 10  
C 1  
D  $\frac{1}{100}$
- 14 Which one of the following statements is **not** true about the control rods used in a nuclear reactor?
- A They must absorb neutrons.  
B They must slow down neutrons to thermal speeds.  
C They must retain their shape at high temperatures.  
D The length of rod in the reactor must be variable.



- 15 The magnetic flux,  $\Phi$ , through a coil varies with time,  $t$ , as shown by the first graph. Which one of the following graphs, **A** to **D**, best represents how the magnitude,  $\varepsilon$ , of the induced emf varies in this same period of time?



**END OF SECTION A**

**THERE ARE NO QUESTIONS PRINTED ON THIS PAGE**

**THERE ARE NO QUESTIONS PRINTED ON THIS PAGE**

**THERE ARE NO QUESTIONS PRINTED ON THIS PAGE**